

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
3 November 2005 (03.11.2005)

PCT

(10) International Publication Number  
**WO 2005/103765 A1**

(51) International Patent Classification<sup>7</sup>: **G01V 1/36**

[GB/AU]; 47 Boscombe Avenue, City Beach, W.A. 6015 (AU). **BISLEY, Richard** [GB/AU]; 5/3 Thirlmere Road, Mount Lawley, W.A. 6050 (AU).

(21) International Application Number:  
PCT/US2004/023120

(22) International Filing Date: 16 July 2004 (16.07.2004)

(74) Agent: **PATTERSON, William, B.**; Moser, Patterson & Sheridan, L.L.P., 3040 Post Oak Boulevard, Suite 1500, Houston, TX 77056 (US).

(25) Filing Language: English

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(26) Publication Language: English

(30) Priority Data:  
60/560,223 7 April 2004 (07.04.2004) US

(71) Applicant (for all designated States except US): **WEST-ERNGECO, L.L.C.** [US/US]; 10001 Richmond Avenue, Houston, TX 77042 (US).


(72) Inventors; and

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH,

(75) Inventors/Applicants (for US only): **MOORE, Ian**

[Continued on next page]

(54) Title: FAST 3-D SURFACE MULTIPLE PREDICTION



(57) Abstract: A method and apparatus for predicting a plurality of surface multiples for a plurality of traces in a record of seismic data. In one embodiment, the method includes providing a plurality of target traces at a nominal offset and a nominal azimuth; selecting a plurality of pairs of input traces, wherein the midpoints of the input traces in each pair are separated by half the nominal offset and the azimuth of a line connecting the midpoints of the input traces in each pair is equal to the nominal azimuth; convolving the selected pairs of input traces to generate a plurality of convolutions; and applying a three dimensional operator to the convolutions.

WO 2005/103765 A1



GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— with international search report

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*